Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 1998
Springer-Verlag GmbH & Co.KG, D-14197 Berlin, Germany

Contents

Full instructions to authors

A 6

Editorial

Waldhäusl W.: Where do we want to go? 1

Review

Cherrington A. D., Edgerton D., Sindelar D. K.: The direct and indirect effects of insulin on hepatic glucose production in vivo 987

Efrat S.: Prospects for gene therapy of insulin-dependent diabetes mellitus 1401

Parving H.-H.: Renoprotection in diabetes: genetic and non-genetic risk factors and treatment 745

Porte D. Jr., Seeley R. J., Woods S. C., Baskin D. G., Figlewicz D. P., Schwartz M. W.: Obesity, diabetes and the central nervous system 863

Trovati M., Anfossi G.: Insulin, insulin resistance and platelet function: similarities with insulin effects on cultured vascular smooth muscle cells 609

Yki-Järvinen H., Utriainen T.: Insulin-induced vasodilatation: physiology or pharmacology? 369

Originals

Ader M., Richey J. M., Bergman R. N.: Evidence for direct action of alloxan to induce insulin resistance at the cellular level 1327

Ahrén B., Oosterwijk C., Lips C. J. M., Höppener J. W. M.: Transgenic overexpression of human islet amyloid polypeptide inhibits insulin secretion and glucose elimination after gastric glucose gavage in mice 1374

Almind K., Ambye L., Urhammer S. A., Hansen T., Echwald S. M., Holst J. J., Gromada J., Thorens B., Pedersen O.: Discovery of amino acid variants in the human glucose-dependent insulinotropic polypeptide (GIP) receptor: the impact on the pancreatic beta cell responses and functional expression studies in Chinese hamster fibroblast cells 1194

Anastasiou E., Alevizaki M., Grigorakis S. J., Philippou G., Kyprianou M., Souvatzoglou A.: Decreased stature in gestational diabetes mellitus 997

Augstein P., Elefanty A. G., Allison J., Harrison L. C.: Apoptosis and beta-cell destruction in pancreatic islets of NOD mice with spontaneous and cyclophosphamide-accelerated diabetes 1381

Bakker S. J. L., Hoogeveen E. K., Nijpels G., Kostense P. J., Dekker J. M., Gans R. O. B., Heine R. J.: The association of dietary fibres with glucose tolerance is partly explained by concomitant intake of thiamine: The Hoorn Study 1168

Bannan S., Mansfield M. W., Grant P. J.: Soluble vascular cell adhesion molecule-1 and E-selectin levels in relation to vascular risk factors and to E-selectin genotype in the first degree relatives of NIDDM patients and in NIDDM patients 460

Barkai L., Vámosi I., Lukács K.: Prospective assessment of severe hypoglycaemia in diabetic children and adolescents with impaired and normal awareness of hypoglycaemia 898

Bazzan M., Gruden G., Stella S., Vaccarino A., Tamponi G., Olivetti C., Giunti S., Cavallo-Perin P.: Microalbuminuria in IDDM is associated with increased expression of monocyte procoagulant activity 767

Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 1998
Springer-Verlag GmbH & Co.KG, D-14197 Berlin, Germany

Contents

Full instructions to authors

A 6

Editorial

Waldhäusl W.: Where do we want to go? 1

Review

Cherrington A. D., Edgerton D., Sindelar D. K.: The direct and indirect effects of insulin on hepatic glucose production in vivo 987

Efrat S.: Prospects for gene therapy of insulin-dependent diabetes mellitus 1401

Parving H.-H.: Renoprotection in diabetes: genetic and non-genetic risk factors and treatment 745

Porte D. Jr., Seeley R. J., Woods S. C., Baskin D. G., Figlewicz D. P., Schwartz M. W.: Obesity, diabetes and the central nervous system 863

Trovati M., Anfossi G.: Insulin, insulin resistance and platelet function: similarities with insulin effects on cultured vascular smooth muscle cells 609

Yki-Järvinen H., Utriainen T.: Insulin-induced vasodilatation: physiology or pharmacology? 369

Originals

Ader M., Richey J. M., Bergman R. N.: Evidence for direct action of alloxan to induce insulin resistance at the cellular level 1327

Ahrén B., Oosterwijk C., Lips C. J. M., Höppener J. W. M.: Transgenic overexpression of human islet amyloid polypeptide inhibits insulin secretion and glucose elimination after gastric glucose gavage in mice 1374

Almind K., Ambye L., Urhammer S. A., Hansen T., Echwald S. M., Holst J. J., Gromada J., Thorens B., Pedersen O.: Discovery of amino acid variants in the human glucose-dependent insulinotropic polypeptide (GIP) receptor: the impact on the pancreatic beta cell responses and functional expression studies in Chinese hamster fibroblast cells 1194

Anastasiou E., Alevizaki M., Grigorakis S. J., Philippou G., Kyprianou M., Souvatzoglou A.: Decreased stature in gestational diabetes mellitus 997

Augstein P., Elefanty A. G., Allison J., Harrison L. C.: Apoptosis and beta-cell destruction in pancreatic islets of NOD mice with spontaneous and cyclophosphamide-accelerated diabetes 1381

Bakker S. J. L., Hoogeveen E. K., Nijpels G., Kostense P. J., Dekker J. M., Gans R. O. B., Heine R. J.: The association of dietary fibres with glucose tolerance is partly explained by concomitant intake of thiamine: The Hoorn Study 1168

Bannan S., Mansfield M. W., Grant P. J.: Soluble vascular cell adhesion molecule-1 and E-selectin levels in relation to vascular risk factors and to E-selectin genotype in the first degree relatives of NIDDM patients and in NIDDM patients 460

Barkai L., Vámosi I., Lukács K.: Prospective assessment of severe hypoglycaemia in diabetic children and adolescents with impaired and normal awareness of hypoglycaemia 898

Bazzan M., Gruden G., Stella S., Vaccarino A., Tamponi G., Olivetti C., Giunti S., Cavallo-Perin P.: Microalbuminuria in IDDM is associated with increased expression of monocyte procoagulant activity 767

Benhamou P. Y., Moriscot C., Richard M. J., Beatrix O., Badet L., Pattou F., Kerr-Conte J., Chroboczek J., Lemarchand P., Halimi S.: Adenovirus-mediated catalase gene transfer reduces oxidant stress in human, porcine and rat pancreatic islets 1093

Berg U. B., Torbjörnsdotter T. B., Jaremko G., Thalme B.: Kidney morphological changes in relation to long-term renal function and metabolic control in adolescents with IDDM 1047

Bernardi M., La Rocca E., Castoldi R., Di Carlo V., Caldara R., Furiani S., Giudici D., Pozza G., Secchi A.: Monooligoclonal immunoglobulin abnormalities in diabetic patients after kidney transplantation: influence of simultaneous pancreas graft 1176

Bouwens L., Pipeleers D. G.: Extra-insular beta cells associated with ductules are frequent in adult human pancreas 629

Brod S. A., Malone M., Darcan S., Papolla M., Nelson L.: Ingested interferon α suppresses Type I diabetes in non-obese diabetic mice 1227

Brown G. R., Silva M. D., Thompson P. A., Beutler B.: Lymphoid Hyperplasia, $CD45RB^{high}$ to $CD45RB^{low}$ T-cell imbalance, and suppression of Type I diabetes mellitus result from TNF blockade in NOD → NOD-*scid* adoptive T cell transfer 1502

Buyken A. E., Toeller M., Heitkamp G., Vitelli F., Stehle P., Scherbaum W. A., Fuller J. H., EURODIAB IDDM Complications Study Group.: Relation of fibre intake to HbA_{1c} and the prevalence of severe ketoacidosis and severe hypoglycaemia 882

Cabezas-Cerrato J for the Neuropathy Spanish Study Group of the Spanish Diabetes Society (S. D. S): The prevalence of clinical diabetic polyneuropathy in Spain: a study in primary care and hospital clinic groups 1263

Cameron N. E., Cotter M. A., Horrobin D. H., Tritschler H. J.: Effects of α-lipoic acid on neurovascular function in diabetic rats: interaction with essential fatty acids 390

Carlsson P.-O., Berne C., Jansson L.: Angiotensin II and the endocrine pancreas: effects on islet blood flow and insulin secretion in rats 127

Ceriello A., Mercuri F., Fabbro D., Giacomello R., Stel G., Taboga C., Tonutti L., Motz E., Damante G.: Effect of intensive glycaemic control on fibrinogen plasma concentrations in patients with Type II diabetes mellitus. Relation with β-fibrinogen genotype 1270

Chèvre J.-C., Hani E. H., Boutin P., Vaxillaire M., Blanché H., Vionnet N., Pardini V. C., Timsit J., Larger E., Charpentier G., Beckers D., Maes M., Bellanné-Chantelot C., Velho G., Froguel P.: Mutation screening in 18 Caucasian families suggest the existence of other MODY genes 1017

Chico A., Pérez A., Córdoba A., Arcelú R., Carreras G., Leiva A de., González-Sastre F., Blanco-Vaca F.: Plasma homocysteine is related to albumin excretion rate in patients with diabetes mellitus: a new link between diabetic nephropathy and cardiovascular disease? 684

Cinti S., Eberbach S., Castellucci M., Accili D.: Lack of insulin receptors affects the formation of white adipose tissue in mice. A morphometric and ultrastructural analysis 171

Coutant R., Landais P., Rosilio M., Johnsen C., Lahlou N., Chatelain P., Carel J. C., Ludvigsson J., Boitard C., Bougnères P. F.: Low dose linomide in Type I juvenile diabetes of recent onset: a randomised placebo-controlled double blind trial 1040

Crowther N. J., Cameron N., Trusler J., Gray I. P.: Association between poor glucose tolerance and rapid post natal weight gain in seven-year-old children 1163

Dabelea D., Hanson R. L., Bennett P. H., Roumain J., Knowler W. C., Pettitt D. J.: Increasing prevalence of Type II diabetes in American Indian children 904

Darville M. I., Eizirik D. L.: Regulation by cytokines of the inducible nitric oxide synthase promoter in insulin-producing cells 1101

Day C. P., Grove J., Daly A. K., Stewart M. W., Avery P. J., Walker M.: Tumour necrosis factor-alpha gene promoter polymorphism and decreased insulin resistance 430

Deacon C. F., Knudsen L. B., Madsen K., Wiberg F. C., Jacobsen O., Holst J. J.: Dipeptidyl peptidase IV resistant analogues of glucagon-like peptide-1 which have extended metabolic stability and improved biological activity 271

Derylo B., Babazono T., Glogowski E., Kapor-Drezgic J., Hohman T., Whiteside C.: High glucose-induced mesangial cell altered contractility: role of the polyol pathway 507

Detaille D., Wiernsperger N., Devos P.: Potentiating effect of metformin on insulin-induced glucose uptake and glycogen metabolism within *Xenopus* oocytes 2

Diraison F., Large V., Brunengraber H., Beylot M.: Non-invasive tracing of liver intermediary metabolism in normal subjects and in moderately hyperglycaemic NIDDM subjects. Evidence against increased gluconeogenesis and hepatic fatty acid oxidation in NIDDM 212

Du X. L., Sui G. Z., Stockklauser-Färber K., Weiß J., Zink S., Schwippert B., Wu Q. X., Tschöpe D., Rösen P.: Induction of apoptosis by high proinsulin and glucose in cultured human umbilical vein endothelial cells is mediated by reactive oxygen species 249

Dufayet De La Tour D., Raccah D., Jannot M. F., Coste T., Rougerie C., Vague P.: Erythrocyte Na/K ATPase activity and diabetes: relationship with C-peptide level 1080

Ebeling P., Essén-Gustavsson B., Tuominen J. A., Koivisto V. A.: Intramuscular triglyceride content is increased in IDDM 111

Enoksson S., Degerman E., Hagström-Toft E., Large V., Arner P.: Various phosphodiesterase subtypes mediate the in vivo antilipolytic effect of insulin on adipose tissue and skeletal muscle in man 560

Eriksson K.-F., Lindgärde F.: No excess 12-year mortality in men with impaired glucose tolerance who participated in the Malmö Preventive Trial with diet and exercise 1010

Eriksson P., Reynisdottir S., Lönnqvist F., Stemme V., Hamsten A., Arner P.: Adipose tissue secretion of plasminogen activator inhibitor-1 in non-obese and obese individuals 65

Fagot-Campagna A., Nelson R. G., Knowler W. C., Pettitt D. J., Robbins D. C., Go O., Welty T. K., Lee E. T., Howard B. V.: Plasma lipoproteins and the incidence of abnormal excretion of albumin in diabetic American Indians: The Strong Heart Study 1002

Fernyhough P., Diemel L. T., Tomlinson D. R.: Target tissue production and axonal transport of neurotrophin-3 are reduced in streptozotocin-diabetic rats 300

Festa A., Kopp H. P., Schernthaner G., Menzel E. J.: Autoantibodies to oxidised low density lipoproteins in IDDM are inversely related to metabolic control and microvascular complications 350

Festa A., Schmölzer B., Schernthaner G., Menzel E. J.: Differential expression of receptors for advanced glycation end products on monocytes in patients with IDDM 674

Fogarty D. G., Zychma M. J., Scott L. J., Warram J. H., Krolewski A. S.: The C825T polymorphism in the human G-protein $\beta 3$ subunit gene is not associated with diabetic nephropathy in Type I diabetes mellitus 1304

Forsblom C. M., Sane T., Groop P.-H., Tötterman K. J., Kallio M., Saloranta C., Laasonen L., Summanen P., Lepäntalo M., Laatikainen L., Matikainen E., Teppo A.-M., Koskimies S., Groop L.: Risk factors for mortality in Type II (non-insulin-dependent) dia-

betes: evidence of a role for neuropathy and a protective effect of HLA-DR4 1253
Frystyk J., Skjærbæk C., Zapf J., Ørskov H.: Increased levels of circulating free insulin-like growth factors in patients with non-islet cell tumour hypoglycaemia 589
Fu J., Ikegami H., Kawaguchi Y., Fujisawa T., Kawabata Y., Hamada Y., Ueda H., Shintani M., Nojima K., Babaya N., Shen Q.-J., Uchigata Y., Urakami T., Omori Y., Shima K., Ogihara T.: Association of distal chromosome 2q with IDDM in Japanese subjects 228
Füchtenbusch M., Rabl W., Grassl B., Bachmann W., Standl E., Ziegler A.-G.: Delay of Type I diabetes in high risk, first degree relatives by parenteral antigen administration: the Schwabing Insulin Prophylaxis Pilot Trial 536
Fujisawa T., Ikegami H., Kawaguchi Y., Hamada Y., Ueda H., Shintani M., Fukuda M., Ogihara T.: Meta-analysis of association of insertion/deletion polymorphism of angiotensin I-converting enzyme gene with diabetic nephropathy and retinopathy 47
Fujita Y., Kojima H., Hidaka H., Fujimiya M., Kashiwagi A., Kikkawa R.: Increased intestinal glucose absorption and postprandial hyperglycaemia at the early step of glucose intolerance in Otsuka Long-Evans Tokushima Fatty Rats 1459
Fürnsinn C., Brunmair B., Furtmüller R., Roden M., Englisch R., Waldhäusl W.: Failure of leptin to affect basal and insulin-stimulated glucose metabolism of rat skeletal muscle in vitro 524
Garofano A., Czernichow P., Bréant B.: Beta-cell mass and proliferation following late fetal and early postnatal malnutrition in the rat 1114
Giaccari A., Morviducci L., Pastore L., Zorretta D., Sbraccia P., Maroccia E., Buongiorno A., Tamburrano G.: Relative contribution of glycogenolysis and gluconeogenesis to hepatic glucose production in control and diabetic rats. A re-examination in the presence of euglycaemia 307
Gibbins R. L., Owens D. R., Allen J. C., Eastman L.: Practical application of the European Field Guide in screening for diabetic retinopathy by using ophthalmoscopy and 35 mm retinal slides 59
Guenifi A., Abdel-Halim S. M., Efendic S., Östenson C.-G.: Preserved initiatory and potentiatory effect of α-ketoisocaproate on insulin release in islets of glucose intolerant rats 1368
Hahl J., Simell T., Ilonen J., Knip M., Simell O.: Costs of predicting IDDM 79
Hajduch E., Darakhshan F., Hundal H. S.: Fructose uptake in rat adipocytes: GLUT5 expression and the effects of streptozotocin-induced diabetes 821
Hallakou S., Foufelle F., Doaré L., Kergoat M., Ferré P.: Pioglitazone-induced increase of insulin sensitivity in the muscles of the obese Zucker fa/fa rat cannot be explained by local adipocyte differentiation 963
Hammes H.-P., Wellensiek B., Klöting I., Sickel E., Bretzel R. G., Brownlee M.: The relationship of glycaemic level to advanced glycation end-product (AGE) accumulation and retinal pathology in the spontaneous diabetic hamster 165
Hampson G., Evans C., Petitt R. J., Evans W. D., Woodhead S. J., Peters J. R., Ralston S. H.: Bone mineral density, collagen type 1 α 1 genotypes and bone turnover in premenopausal women with diabetes mellitus 1314
Hani E. H., Boutin P., Durand E., Inoue H., Permutt M. A., Velho G., Froguel P.: Missense mutations in the pancreatic islet beta cell inwardly rectifying K^+ channel gene (*KIR6.2/BIR*): a meta-analysis suggests a role in the polygenic basis of Type II diabetes mellitus in Caucasians 1511
Hirata K., Shikata K., Matsuda M., Akiyama K., Sugimoto H., Kushiro M., Makino H.: Increased expression of selectins in kidneys of patients with diabetic nephropathy 185
Holl R. W., Grabert M., Sorgo W., Heinze E., Debatin K. M.: Contributions of age, gender and insulin administration to weight gain in subjects with IDDM 542
Hopfner R. L., Misurski D., Wilson T. W., McNeill J. R., Gopalakrishnan V.: Insulin and vanadate restore decreased plasma endothelin concentrations and exaggerated vascular responses to normal in the streptozotocin diabetic rat 1233
Hovorka R., Koukkou E., Southerden D., Powrie J. K., Young M. A.: Measuring prehepatic insulin secretion using a population model of C-peptide kinetics: accuracy and required sampling schedule 548
Hudson C., Flanagan J. G., Turner G. S., Chen H. C., Young L. B., McLeod D.: Influence of laser photocoagulation for clinically significant diabetic macular oedema (DMO) on short-wavelength and conventional automated perimetry 1283
Hudson C., Flanagan J. G., Turner G. S., Chen H. C., Young L. B., McLeod D.: Short-wavelength sensitive visual field loss in patients with clinically significant diabetic macular oedema 918
Hussain M. J., Maher J., Warnock T., Vats A., Peakman M., Vergani D.: Cytokine overproduction in healthy first degree relatives of patients with IDDM 343
Ibáñez L., Potau N., Chacon P., Pascual C., Carrascosa A.: Hyperinsulinaemia, dyslipaemia and cardiovascular risk in girls with a history of premature pubarche 1057
Inada H., Shindo H., Tawata M., Onaya T.: cAMP regulates nitric oxide production and ouabain sensitive Na^+, K^+-ATPase activity in SH-SY5Y human neuroblastoma cells 1451
Ishiyama N., Kanzaki M., Seno M., Yamada H., Kobayashi I., Kojima I.: Studies on the betacellulin receptor in pancreatic AR42J cells 623
Isomaa B., Henricsson M., Lehto M., Forsblom C., Karanko S., Sarelin L., Häggblom M., Groop L.: Chronic diabetic complications in patients with MODY3 diabetes 467
Jager A., Kostense P. J., Nijpels G., Heine R. J., Bouter L. M., Stehouwer C. D. A.: Microalbuminuria is strongly associated with NIDDM and hypertension, but not with the insulin resistance syndrome: the Hoorn Study 694
Jang H. C., Min H. K., Lee H. K., Cho N. H., Metzger B. E.: Short stature in Korean women: a contribution to the multifactorial predisposition to gestational diabetes mellitus 778
Jones S. C., Thomas T. H., Marshall S. M.: Abnormal regulation of cell membrane fluidity in diabetic nephropathy 337
Jörneskog G., Brismar K., Fagrell B.: Pronounced skin capillary ischemia in the feet of diabetic patients with bad metabolic control 410
Jude E. B., Abbott C. A., Young M. J., Anderson S. G., Douglas J. T., Boulton A. J. M.: The potential role of cell adhesion molecules in the pathogenesis of diabetic neuropathy 330
Kanc K., Janssen M. M. J., Keulen E. T. P., Jacobs M. A. J. M., Popp-Snijders C., Snoek F. J., Heine R. J.: Substitution of night-time continuous subcutaneous insulin infusion therapy for bedtime NPH insulin in a multiple injection regimen improves counterregulatory hormonal responses and warning symptoms of hypoglycaemia in IDDM 322
Kawamura T., Umemura T., Kanai A., Uno T., Matsumae H., Sano T., Sakamoto N., Sakakibara T., Nakamura J., Hotta N.: The incidence and characteristics of silent cerebral infarction in elderly diabetic patients: association with serum-soluble adhesion molecules 911
Keymeulen B., Ling Z., Gorus F. K., Delvaux G., Bouwens L., Grupping A., Hendrieckx C., Pipeleers-Marichal M., Van Schravendijk C., Salmela K., Pipeleers D. G.: Implantation of standardized beta-cell grafts in a liver segment of IDDM patients: graft and recipient

characteristics in two cases of insulin-independence under maintenance immunosuppression for prior kidney graft 452
Klannemark M., Orho M., Langin D., Laurell H., Holm C., Reynisdottir S., Arner P., Groop L.: The putative role of the hormone-sensitive lipase gene in the pathogenesis of Type II diabetes mellitus and abdominal obesity 1516
Kong M.-F., King P., Macdonald I. A., Blackshaw P. E., Perkins A. C., Armstrong E., Buchanan K. D., Tattersall R. B.: Effect of euglycaemic hyperinsulinaemia on gastric emptying and gastrointestinal hormone responses in normal subjects 474
Kong M.-F., Stubbs T. A., King P., Macdonald I. A., Lambourne J. E., Blackshaw P. E., Perkins A. C., Tattersall R. B.: The effect of single doses of pramlintide on gastric emptying of two meals in men with IDDM 577
Koopmans S. J., Leighton B., DeFronzo R. A.: Neonatal de-afferentation of capsaicin-sensitive sensory nerves increases in vivo insulin sensitivity in conscious adult rats 813
Koren W., Koldanov R., Pronin V. S., Postnov I. Y., Peleg E., Rosenthal T., Berezin M., Postnov Y. V.: Enhanced erythrocyte Na^+/H^+ exchange predicts diabetic nephropathy in patients with IDDM 201
Kuzuya M., Satake S., Ai S., Asai T., Kanda S., Ramos M. A., Miura H., Ueda M., Iguchi A.: Inhibition of angiogenesis on glycated collagen lattices 491
Lam K. S. L., Ma O. C. K., Bourke C., Chan L. C., Janus E. D.: Genetic influence of the R/Q353 genotype on factor VII activity is overwhelmed by environmental factors in Chinese patients with Type II (non-insulin-dependent) diabetes mellitus 760
Lamhamedi-Cherradi S.-E., Luan J.-J., Eloy L., Fluteau G., Bach J.-F., Garchon H.-J.: Resistance of T-cells to apoptosis in autoimmune diabetic (NOD) mice is increased early in life and is associated with dysregulated expression of Bcl-x 178
Larsson H., Ahrén B.: Glucose-dependent arginine stimulation test for characterization of islet function: studies on reproducibility and priming effect of arginine 772
LeBras S., Czernichow P., Scharfmann R.: A search for tyrosine kinase receptors expressed in the rat embryonic pancreas 1474
Lindsay R. M., Jamieson N. S. D., Walker S. A., McGuigan C. C., Smith W., Baird J. D.: Tissue ascorbic acid and polyol pathway metabolism in experimental diabetes 516
Ling Z.-C., Khan A., Delauny F., Davani B., Östenson C.-G., Gustafsson J.-Å., Okret S., Landau B. R., Efendic S.: Increased glucocorticoid sensitivity in islet beta-cells: effects on glucose 6-phosphatase, glucose cycling and insulin release 634
Lorenzen T., Pociot F., Stilgren L., Kristiansen O. P., Johannesen J., Olsen P. B., Walmar A., Larsen A., Albrechtsen N. C., Eskildsen P. C., Andersen O. O., Nerup J and the Danish IDDM Epidemiology and Genetics Group.: Predictors of IDDM recurrence risk in offspring of Danish IDDM patients 666
Makino A., Kamata K.: Possible modulation by endothelin-1, nitric oxide, prostaglandin I_2 and thromboxane A_2 of vasoconstriction induced by an α-agonist in mesenteric arterial bed from diabetic rats 1410
Malkki M., Laakso M., Deeb S. S.: Functional consequences of naturally occurring variants of human hexokinase II 1205
Mari A.: Assessment of insulin sensitivity and secretion with the labelled intravenous glucose tolerance test: improved modelling analysis 1029
Martin S., Heidenthal E., Schulte B., Rothe H., Kolb H.: Soluble forms of intercellular adhesion molecule-1 inhibit insulitis and onset of autoimmune diabetes 1298
McKeigue P. M., Lithell H. O., Leon D. A.: Glucose tolerance and resistance to insulin-stimulated glucose uptake in men aged 70 years in relation to size at birth 1133
Morgan D. G., Kulkarni R. N., Hurley J. D., Wang Z. L., Wang R. M., Ghatei M. A., Karlsen A. E., Bloom S. R., Smith D. M.: Inhibition of glucose stimulated insulin secretion by neuropeptide Y is mediated via the Y1 receptor and inhibition of adenylyl cyclase in RIN 5AH rat insulinoma cells 1482
Motoyoshi S., Shirotani T., Araki E., Sakai K., Kaneko K., Motoshima H., Yoshizato K., Shirakami A., Kishikawa H., Shichiri M.: Cellular characterization of pituitary adenoma cell line (AtT20 cell) transfected with insulin, glucose transporter type 2 (GLUT2) and glucokinase genes: insulin secretion in response to physiological concentrations of glucose 1492
Mühlhauser I., Overmann H., Bender R., Bott U., Berger M.: Risk factors of severe hypoglycaemia in adult patients with Type I diabetes – a prospective population based study 1274
Mühlhauser I., Overmann H., Bender R., Bott U., Jörgens V., Trautner Ch., Siegrist J., Berger M.: Social status and the quality of care for adult people with Type I (insulin-dependent) diabetes mellitus – a population-based study 1139
Murase K., Odaka H., Suzuki M., Tayuki N., Ikeda H.: Pioglitazone time-dependently reduces tumour necrosis factor-α level in muscle and improves metabolic abnormalities in Wistar fatty rats 257
Murata T., Hoffmann S., Ishibashi T., Spee C., Gordon E. M., Anderson W. F., Hinton D. R., Ryan S. J.: Retrovirus-mediated gene transfer targeted to retinal photocoagulation sites 500
Nakazaki M., Ishihara H., Kakei M., Inukai K., Asano T., Miyazaki J.-I., Tanaka H., Kikuchi M., Yada T., Oka Y.: Repetitive mitochondrial Ca^{2+} signals synchronize with cytosolic Ca^{2+} oscillations in the pancreatic beta-cell line, MIN6 279
Naserke H. E., Dozio N., Ziegler A.-G., Bonifacio E.: Comparison of a novel micro-assay for insulin autoantibodies with the conventional radiobinding assay 681
New J. P., Marshall S. M., Bilous R. W.: Renal autoregulation is normal in newly diagnosed, normotensive, NIDDM patients 206
Ng L. L., Ngkeekwong F. C., Quinn P. A., Davies J. E.: Uptake mechanisms for ascorbate and dehydroascorbate in lymphoblasts from diabetic nephropathy and hypertensive patients 435
Niculescu L., Van Schaftingen E.: Mannitol 1-phosphate mediates an inhibitory effect of mannitol on the activity and the translocation of glucokinase in isolated rat hepatocytes 947
Nielsen M. F., Dinneen S., Basu A., Basu R., Alzaid A., Rizza R. R.: Failure of nocturnal changes in growth hormone to alter carbohydrate tolerance the following morning 1064
Niklasson M., Holmäng A., Lönnroth P.: Induction of rat muscle insulin resistance by epinephrine is accompanied by increased interstitial glucose and lactate concentrations 1467
Nordfors L., Hoffstedt J., Nyberg B., Thörne A., Arner P., Schalling M., Lönnqvist F.: Reduced gene expression of UCP2 but not UCP3 in skeletal muscle of human obese subjects 935
Oberkofler H., Liu Y. M., Esterbauer H., Hell E., Krempler F., Patsch W.: Uncoupling protein-2 gene: reduced mRNA expression in intraperitoneal adipose tissue of obese humans 940
Obrosova I., Cao X., Greene D. A., Stevens M. J.: Diabetes-induced changes in lens antioxidant status, glucose utilization and energy metabolism: effect of DL-α-lipoic acid 1442
O'Harte F. P. M., Abdel-Wahab Y. H. A., Conlon J. M., Flatt P. R.: Glycation of glucagon-like peptide-1(7–36)amide:

characterization and impaired action on rat insulin secreting cells 1187
Ohtomo Y., Bergman T., Johansson B.-L., Jörnvall H., Wahren J.: Differential effects of proinsulin C-peptide fragments on Na^+, K^+-ATPase activity of renal tubule segments 287
Ono Y., Umeda F., Kunisaki M., Sekiguchi N., Hashimoto T., Nawata H.: Effect of high glucose concentrations on prostacyclin-stimulating factor mRNA expression in cultured aortic smooth muscle cells 134
Ozanne S. E., Martensz N. D., Petry C. J., Loizou C. L., Hales C. N.: Maternal low protein diet in rats programmes fatty acid desaturase activities in the offspring 1337
Pagano P. J., Griswold M. C., Ravel D., Cohen R. A.: Vascular action of the hypoglycaemic agent gliclazide in diabetic rabbits 9
Palmer A. M., Gopaul N., Dhir S., Thomas C. R., Poston L., Tribe R. M.: Endothelial dysfunction in streptozotocin-diabetic rats is not reversed by dietary probucol or simvastatin supplementation 157
Palmer A. M., Thomas C. R., Gopaul N., Dhir S., Änggård E. E., Poston L., Tribe R. M.: Dietary antioxidant supplementation reduces lipid peroxidation but impairs vascular function in small mesenteric arteries of the streptozotocin-diabetic rat 148
Paolisso G., Tagliamonte M. R., Rizzo M. R., Gualdiero P., Saccomanno F., Gambardella A., Giugliano D., D'Onofrio F., Howard B. V.: Lowering fatty acids potentiates acute insulin response in first degree relatives of people with Type II diabetes 1127
Peak M., Rochford J. J., Borthwick A. C., Yeaman S. J., Agius L.: Signalling pathways involved in the stimulation of glycogen synthesis by insulin in rat hepatocytes 16
Peschke E., Peschke D.: Evidence for a circadian rhythm of insulin release from perifused rat pancreatic islets 1085
Petersen J. S., Rimvall K., Jørgensen P. N., Hasselager E., Moody A., Hejnæs K., Clausen J. T., Dyrberg T.: Regulation of GAD expression in rat pancreatic islets and brain by γ-vinyl-GABA and glucose 530
Pieper G. M., Dembny K., Siebeneich W.: Long-term treatment in vivo with NOX-101, a scavenger of nitric oxide, prevents diabetes-induced endothelial dysfunction 1220
Pieper G. M., Dondlinger L. A.: Antioxidant pyrrolidine dithiocarbamate prevents defective bradykinin-stimulated calcium accumulation and nitric oxide activity following exposure of endothelial cells to elevated glucose concentration 806
Pietruck F., Spleiter S., Daul A., Philipp T., Derwahl M., Schatz H., Siffert W.: Enhanced G protein activation in IDDM patients with diabetic nephropathy 94
Pijl JW van der., Daha M. R., Born J van den., Verhagen N. A. M., Lemkes H. H. P. J., Bucala R., Berden J. H. M., Zwinderman A. H., Bruijn J. A., Es LA van., Woude FJ van der.: Extracellular matrix in human diabetic nephropathy: reduced expression of heparan sulphate in skin basement membrane 791
Poulsen P. L., Bek T., Ebbehøj E., Hansen K. W., Mogensen C. E.: 24-h ambulatory blood pressure and retinopathy in normoalbuminuric IDDM patients 105
Poulton J., Scott-Brown M., Cooper A., Marchington D. R., Phillips D. I. W.: A common mitochondrial DNA variant is associated with insulin resistance in adult life 54
Qiang X., Satoh J., Sagara M., Fukuzawa M., Masuda T., Sakata Y., Muto G., Muto Y., Takahashi K., Toyota T.: Inhibitory effect of troglitazone on diabetic neuropathy in streptozotocin-induced diabetic rats 1321
Raabe H.-M., Höpner J.-H., Notbohm H., Sinnecker G. H. G., Kruse K., Müller P. K.: Biochemical and biophysical alterations of the 7S and NC1 domain of collagen IV from human diabetic kidneys 1073
Rabini R. A., Staffolani R., Fumelli P., Mutus B., Curatola G., Mazzanti L.: Decreased nitric oxide synthase activity in platelets from IDDM and NIDDM patients 101
Ranganath L., Sedgwick I., Morgan L., Wright J., Marks V.: The ageing entero-insular axis 1309
Renaudin C., Michoud E., Rapin J. R., Lagarde M., Wiernsperger N.: Hyperglycaemia modifies the reaction of microvessels to insulin in rat skeletal muscle 26
Riemens S. C., van Tol A., Sluiter W. J., Dullaart R. P. F.: Plasma phospholipid transfer protein activity is related to insulin resistance: impaired acute lowering by insulin in obese Type II diabetic patients 929
Romano G., Tilly-Kiesi M. K., Patti L., Taskinen M.-R., Pacioni D., Cassader M., Riccardi G., Rivellese A. A.: Effects of dietary cholesterol on plasma lipoproteins and their subclasses in IDDM patients 193
Roufail E., Soulis T., Boel E., Cooper M. E., Rees S.: Depletion of nitric oxide synthase-containing neurons in the diabetic retina: reversal by aminoguanidine 1419
Russell J. C., Dolphin P. J., Graham S. E., Amy R. M., Brindley D. N.: Improvement of insulin sensitivity and cardiovascular outcomes in the JCR:LA-cp rat by D-fenfluramine 380
Sackmann H., Tran-Van T., Tack I., Hanaire-Broutin H., Tauber J.-P., Ader J.-L.: Renal functional reserve in IDDM patients 86
Sakura H., Ashcroft S. J. H., Terauchi Y., Kadowaki T., Ashcroft F. M.: Glucose modulation of ATP-sensitive K-currents in wild-type, homozygous and heterozygous glucokinase knock-out mice 654
Salbe A. D., Fontvieille A. M., Pettitt D. J., Ravussin E.: Maternal diabetes status does not influence energy expenditure or physical activity in 5-year-old Pima Indian children 1157
Sano T., Umeda F., Hashimoto T., Nawata H., Utsumi H.: Oxidative stress measurement by in vivo electron spin resonance spectroscopy in rats with streptozotocin-induced diabetes 1355
Saukkonen T., Virtanen S. M., Karppinen M., Reijonen H., Ilonen J., Räsänen L., Åkerblom H. K., Savilahti E., and the Childhood Diabetes in Finland Study Group.: Significance of cow's milk protein antibodies as risk factor for childhood IDDM: interactions with dietary cow's milk intake and HLA-DQB1 genotype 72
Savola K., Bonifacio E., Sabbah E., Kulmala P., Vähäsalo P., Karjalainen J., Tuomilehto-Wolf E., Meriläinen J., Åkerblom H. K., Knip M and the Childhood Diabetes in Finland Study Group.: IA-2 antibodies – a sensitive marker of IDDM with clinical onset in childhood and adolescence 424
Savola K., Sabbah E., Kulmala P., Vähäsalo P., Ilonen J., Knip M.: Autoantibodies associated with Type I diabetes mellitus persist after diagnosis in children 1293
Schneider D. J., Absher P. M., Neimane D., Russell J. C., Sobel B. E.: Fibrinolysis and atherogenesis in the JCR:LA-cp rat in relation to insulin and triglyceride concentrations in blood 141
Seissler J., de Sonnaville J. J. J., Morgenthaler N. G., Steinbrenner H., Glawe D., Khoo-Morgenthaler U. Y., Lan M. S., Notkins A. L., Heine R. J., Scherbaum W. A.: Immunological heterogeneity in Type I diabetes: presence of distinct autoantibody patterns in patients with acute onset and slowly progressive disease 891
Sener A., Kadiata M. M., Olivares E., Malaisse W. J.: Comparison of the effects of D-mannoheptulose and its hexaacetate ester on D-glucose metabolism and insulinotropic action in rat pancreatic islets 1109
Serino R., Ueta Y., Tokunaga M., Hara Y., Nomura M., Kabashima N., Shibuya I., Hattori Y., Yamashita H.: Upregulation of hypothalamic nitric oxide

synthase gene expression in streptozotocin-induced diabetic rats 640
Sharpe P. C., Yue K. K. M., Catherwood M. A., McMaster D., Trimble E. R.: The effects of glucose-induced oxidative stress on growth and extracellular matrix gene expression of vascular smooth muscle cells 1210
Songini M., Bernardinelli L., Clayton D., Montomoli C., Pascutto C., Ghislandi M., Fadda D., Bottazzo GF and the Sardinian IDDM Study Group.: The Sardinian IDDM Study: 1. Epidemiology and geographical distribution of IDDM in Sardinia during 1989 to 1994 221
Spinas G. A., Laffranchi R., Francoys I., David I., Richter C., Reinecke M.: The early phase of glucose-stimulated insulin secretion requires nitric oxide 292
Sugimoto H., Shikata K., Matsuda M., Kushiro M., Hayashi Y., Hiragushi K., Wada J., Makino H.: Increased expression of endothelial cell nitric oxide synthase (ecNOS) in afferent and glomerular endothelial cells is involved in glomerular hyperfiltration of diabetic nephropathy 1426
Suzuki S., Hinokio Y., Ohtomo M., Hirai M., Hirai A., Chiba M., Kasuga S., Satoh Y., Akai H., Toyota T.: The effects of coenzyme Q_{10} treatment on maternally inherited diabetes mellitus and deafness, and mitochondrial DNA 3243 (A to G) mutation 584
Sweeney G., Somwar R., Ramlal T., Martin-Vasallo P., Klip A.: Insulin stimulation of K^+ uptake in 3T3-L1 fibroblasts involves phosphatidylinositol 3-kinase and protein kinase C-zeta 1199
Tack C. J. J., Ong M. K. E., Lutterman J. A., Smits P.: Insulin-induced vasodilatation and endothelial function in obesity/insulin resistance. Effects of troglitazone 569
Takeda Y., Gotoh M., Dono K., Nishihara M., Grochowiecki T., Kimura F., Yoshida T., Ohta Y., Ota H., Ohzato H., Umeshita K., Takeda T., Matsuura N., Sakon M., Kayagaki N., Yagita H., Okumura K., Miyasaka M., Monden M.: Protection of islet allografts transplanted together with Fas ligand expressing testicular allografts 315
Temelkova-Kurktschiev T., Koehler C., Schaper F., Henkel E., Hahnefeld A., Fuecker K., Siegert G., Hanefeld M.: Relationship between fasting plasma glucose, atherosclerosis risk factors and carotid intima media thickness in non-diabetic individuals 706
Terasaki J., Anai M., Funaki M., Shibata T., Inukai K., Ogihara T., Ishihara H., Katagiri H., Onishi Y., Sakoda H., Fukushima Y., Yazaki Y., Kikuchi M., Oka Y., Asano T.: Role of JTT-501, a new insulin sensitiser, in restoring impaired GLUT4 translocation in adipocytes of rats fed a high fat diet 400
The Diabetes Control and Complication Trial Research Group.: The effect of intensive diabetes therapy on measures of autonomic nervous system function in the Diabetes Control and Complications Trial (DCCT) 416
The Eurodiab Ace Study Group and The Eurodiab Ace Substudy 2 Study Group.: Familial Risk of Type I Diabetes in European Children 1151
Toft-Nielsen M., Madsbad S., Holst J. J.: Exaggerated secretion of glucagon-like peptide-1 (GLP-1) could cause reactive hypoglycaemia 1180
Traverso N., Menini S., Cosso L., Odetti P., Albano E., Pronzato M. A., Marinari U. M.: Immunological evidence for increased oxidative stress in diabetic rats 265
Tribe R. M., Thomas C. R., Poston L.: Flow-induced dilatation in isolated resistance arteries from control and streptozotocin-diabetic rats 34
Tuomilehto J., Borch-Johnsen K., Molarius A., Forsén T., Rastenyte D., Sarti C., Reunanen A.: Incidence of cardiovascular disease in Type 1 (insulin-dependent) diabetic subjects with and without diabetic nephropathy in Finland 784
Ueda K., Tanizawa Y., Ishihara H., Kizuki N., Ohta Y., Matsutani A., Oka Y.: Overexpression of mitochondrial FAD-linked glycerol-3-phosphate dehydrogenase does not correct glucose-stimulated insulin secretion from diabetic GK rat pancreatic islets 649
Utriainen T., Takala T., Luotolahti M., Rönnemaa T., Laine H., Ruotsalainen U., Haaparanta M., Nuutila P., Yki-Järvinen H.: Insulin resistance characterizes glucose uptake in skeletal muscle but not in the heart in NIDDM 555
Utsunomiya N., Ohagi S., Sanke T., Tatsuta H., Hanabusa T., Nanjo K.: Organization of the human carboxypeptidase E gene and molecular scanning for mutations in Japanese subjects with NIDDM or obesity 701
Valve R., Heikkinen S., Rissanen A., Laakso M., Uusitupa M.: Synergistic effect of polymorphisms in uncoupling protein 1 and β_3-adrenergic receptor genes on basal metabolic rate in obese Finns 357
Vettor R., Pagano C., Granzotto M., Englaro P., Angeli P., Blum W. F., Federspil G., Rohner-Jeanrenaud F., Jeanrenaud B.: Effects of intravenous neuropeptide Y on insulin secretion and insulin sensitivity in skeletal muscle in normal rats 1361
Vreugdenhil G. R., Geluk A., Ottenhoff T. H. M., Melchers W. J. G., Roep B. O., Galama J. M. D.: Molecular mimicry in diabetes mellitus: the homologous domain in coxsackie B virus protein 2C and islet autoantigen GAD_{65} is highly conserved in the coxsackie B-like enteroviruses and binds to the diabetes associated HLA-DR3 molecule 40
Wu S. Q., Kwan C. Y., Tang F.: Streptozotocin-induced diabetes has differential effects on atrial natriuretic peptide synthesis in the rat atrium and ventricle: a study by solution-hybridization-RNase protection assay 660
Yagui K., Shimada F., Mimura M., Hashimoto N., Suzuki Y., Tokuyama Y., Nata K., Tohgo A., Ikehata F., Takasawa S., Okamoto H., Makino H., Saito Y., Kanatsuka A.: A missense mutation in the CD38 gene, a novel factor for insulin secretion: association with Type II diabetes mellitus in Japanese subjects and evidence of abnormal function when expressed in vitro 1024
Yamagishi S., Fujimori H., Yonekura H., Yamamoto Y., Yamamoto H.: Advanced glycation endproducts inhibit prostacyclin production and induce plasminogen activator inhibitor-1 in human microvascular endothelial cells 1435
You S., Gouin E., Saï P.: Spleen cells of non-obese diabetic mice fed with pig splenocytes display modified proliferation and reduced aggressiveness in vitro against pig islet cells 955
Zhu M., Mizuno A., Kuwajima M., Ogino T., Murakami T., Noma Y., Sano T., Shima K.: Ovarian hormone-induced beta-cell hypertrophy contributes to the homeostatic control of beta-cell mass in OLETF female rat, a model of Type II diabetes 799
Ziegler D., Weise F., Langen K.-J., Piolot R., Boy C., Hübinger A., Müller-Gärtner H.-W., Gries F. A.: Effect of glycaemic control on myocardial sympathetic innervation assessed by [^{123}I]metaiodobenzylguanidine scintigraphy: a 4-year prospective study in IDDM patients 443
Zierath J. R., Livingston J. N., Thörne A., Bolinder J., Reynisdottir S., Lönnqvist F., Arner P.: Regional difference in insulin inhibition of non-esterified fatty acid release from human adipocytes: relation to insulin receptor phosphorylation and intracellular signalling through the insulin receptor substrate-1 pathway 1343

Rapid communications

Almind K., Frederiksen S. K., Ahlgren M. G., Urhammer S., Hansen T., Clausen J. O., Pedersen O.: Common amino acid substitutions in insulin receptor substrate-4 are not associated with

Type II diabetes mellitus or insulin resistance 969
Bédard S., Marcotte B., Marette A.: Insulin inhibits inducible nitric oxide synthase in skeletal muscle cells 1523
Bellmann K., Kolb H., Rastegar S., Jee P., Scott F. W.: Potential risk of oral insulin with adjuvant for the prevention of Type I diabetes: a protocol effective in NOD mice may exacerbate disease in BB rats 844
Boeri D., Derchi L. E., Martinoli C., Simoni G., Sampietro L., Storace D., Ponte L., Calvi C., Repetto M., Robaudo C., Maiello M.: Intrarenal arteriosclerosis and impairment of kidney function in NIDDM subjects 121
Ellis T. M., Ottendorfer E., Jodoin E., Salisbury P. J., She J. X., Schatz D. A., Atkinson M. A.: Cellular immune responses to β casein: elevated in but not specific for individuals with Type I diabetes mellitus 731
Fioretto P., Stehouwer C. D. A., Mauer M., Chiesura-Corona M., Brocco E., Carraro A., Bortoloso E., Hinsbergh VWM van., Crepaldi G., Nosadini R.: Heterogeneous nature of microalbuminuria in NIDDM: studies of endothelial function and renal structure 233
Hounsom L., Horrobin D. F., Tritschler H., Corder R., Tomlinson D. R.: A lipoic acid-gamma linolenic acid conjugate is effective against multiple indices of experimental diabetic neuropathy 839
Ishii H., Tada H., Isogai S.: An aldose reductase inhibitor prevents glucose-induced increase in transforming growth factor-β and protein kinase C activity in cultured human mesangial cells 362
Ju Q., Edelstein D., Brendel M. D., Brandhorst D., Brandhorst H., Bretzel R. G., Brownlee M.: Transduction of non-dividing adult human pancreatic beta cells by an integrating lentiviral vector 736
Kellerer M., Mushack J., Seffer E., Mischak H., Ullrich A., Häring H. U.: Protein kinase C isoforms α, δ and θ require insulin receptor substrate-1 to inhibit the tyrosine kinase activity of the insulin receptor in human kidney embryonic cells (HEK 293 cells) 833
Millet L., Vidal H., Larrouy D., Andreelli F., Laville M., Langin D.: mRNA expression of the long and short forms of uncoupling protein-3 in obese and lean humans 829
Radziuk J., Pye S., Bradley B., Braaten J., Vignati L., Roach P., Bowsher R., DiMarchi R., Chance R.: Basal activity profiles of NPH and [N^{ε}-palmitoyl Lys (B29)] human insulins in subjects with IDDM 116
Tian J., Chan C., Kaufman D. L.: Insulin selectively primes Th2 responses and induces regulatory tolerance to insulin in pre-diabetic mice 237
Urhammer S. A., Dalgaard L. T., Sørensen T. I. A., Tybjærg-Hansen A., Echwald S. M., Andersen T., Clausen J. O., Pedersen O.: Organisation of the coding exons and mutational screening of the uncoupling protein 3 gene in subjects with juvenile-onset obesity 241
Urrutia I., Calvo B., Bilbao J. R., the GEPV-N Group., Gastaño L.: Anomalous behaviour of the 5′ insulin gene polymorphism *allele 814:* lack of association with Type I diabetes in Basques 1121
Yamada K., Yuan X., Ishiyama S., Ichikawa F., Kohno S., Shoji S., Hayashi H., Nonaka K.: Identification of a single nucleotide insertion polymorphism in the upstream region of the insulin promoter factor-1 gene: an association study with diabetes mellitus 603

Short communications

De Mattia G., Bravi M. C., Laurenti O., Cassone-Faldetta M., Proietti A., De Luca O., Armiento A., Ferri C.: Reduction of oxidative stress by oral N-acetyl-L-cysteine treatment decreases plasma soluble vascular cell adhesion molecule-1 concentrations in non-obese, non-dyslipidaemic, normotensive, patients with non-insulin-dependent diabetes 1392
Kalidas K., Wasson J., Glaser B., Meyer J. M., Duprat L. J., White M. F., Permutt M. A.: Mapping of the human insulin receptor substrate-2 gene, identification of a linked polymorphic marker and linkage analysis in families with Type II diabetes: no evidence for a major susceptibility role 1389
Møller A. M., Dalgaard L. T., Pociot F., Nerup J., Hansen T., Pedersen O.: Mutations in the hepatocyte nuclear factor-1α gene in Caucasian families originally classified as having Type I diabetes 1528

For debates

Amiel S. A.: Cognitive function testing in studies of acute hypoglycaemia: rights and wrongs? 713
Fanelli C. G., Pampanelli S., Porcellati F., Bolli G. B.: Shift of glycaemic thresholds for cognitive function in hypoglycaemia unawareness in humans 720
Gambaro G., Skrha J., Ceriello A.: Glycosaminoglycan therapy for long-term diabetic complications? 975
Gazis A., Page S. R., Cockcroft J. R.: ACE inhibitors, diabetes and cardiovascular disease 595
Heinemann L., Schmelzeisen-Redeker G on behalf of the Non-invasive task force (N. I. T. F): Non-invasive continuous glucose monitoring in Type I diabetic patients with optical glucose sensors 848
Leahy J. L., Chen Ch., Rhodes Ch. J.: How is the insulin content regulated in nondiabetic hyperinsulinaemic, insulin resistant states? Study of spontaneously hypertensive rats 855
Ozanne S. E., Hales C. N.: Thrifty yes, genetic no 485
Pickup J. C., Crook M. A.: Is Type II diabetes mellitus a disease of the innate immune system? 1241
Reaven G. M.: Hypothesis: muscle insulin resistance is the ("not-so") thrifty genotype 482
Sawicki P. T.: Do ACE inhibitors offer specific benefits in the antihypertensive treatment of diabetic patients? 598
Zierler K.: Does insulin-induced increase in the amount of plasma membrane GLUTs quantitatively account for insulin-induced increase in glucose uptake? 724

Letters to the editors

Baba T., Drenckhan M., Neugebauer S., Klein H.: Effect of angiotensin II and bradykinin on insulin-stimulated tyrosine kinase activity of insulin receptor 741
Bakker S. J. L., Dekker J. M., Heine R. J.: Association between HbA_{1c} and HDL-cholesterol independent of fasting triglycerides in a Caucasian population: evidence for enhanced cholesterol ester transfer induced by in vivo glycation 1249
Bartsocas C. S., Dacou-Voutetakis C., Damianaki D., Karayanni C. H., Kassiou C., Qadreh A., Theodoridis C. H., Tsoka H., Green A., for the Eurodiab ACE G1 Group.: Epidemiology of childhood IDDM in Athens: trends in incidence for the years 1989–1995 245
Berger M., Mühlhauser I., Sawicki P. T.: Possible risk of sulphonylureas in the treatment of non-insulin-dependent diabetes mellitus and coronary artery disease 744
Braun J., Donner H., Plock K., Rau H., Usadel K. H., Badenhoop K.: Hereditary hemochromatosis mutations (HFE) in patients with Type II diabetes mellitus 983
Chuang L.-M., Lee K.-C., Huang C.-N., Wu H.-P., Tai T.-Y., Lin B. J.: Role of S20G mutation of amylin gene in insulin secretion, insulin sensitivity, and Type II diabetes mellitus in Taiwanese patients 1250

Fasching P.: Possible risk of sulphonylureas in the treatment of non-insulin-dependent diabetes mellitus and coronary artery disease 743
Gasic S., Wagner O. F., Fasching P., Ludwig C., Veitl M., Kapiotis S., Jilma B.: Fosinopril decreases levels of soluble vascular cell adhesion molecule-1 in Type II diabetic patients suffering from microalbuminuria 980
Hirai M., Suzuki S., Hinokio Y., Chiba M., Kasuga S., Hirai A., Toyota T.: Group specific component protein genotype is associated with NIDDM in Japan 742
Kaneto H., Ikeda M., Kishimoto M., Iida M., Hoshi A., Watarai T., Kubota M., Kajimoto Y., Yamasaki Y., Hori M.: Dramatic recovery of counter-regulatory hormone response to hypoglycaemia after intensive insulin therapy in poorly controlled Type I diabetes mellitus 982
Katsilambros., Tentolouris N., Linos A., Stambulis E., Papageorgiou K.: Blood pressure and heart rate changes during leaning forward in normal and in NIDDM subjects 126
Kogure A., Yoshida T., Sakane N., Umekawa T., Takakura Y., Kondo M.: Synergic effect of polymorphisms in uncoupling protein 1 and β_3-adrenergic receptor genes on weight loss in obese Japanese 1399
Larsson H., Berglund G., Lindgärde F., Ahrén B.: Comparison of ADA and WHO criteria for diagnosis of diabetes and glucose intolerance 1124
Lobefalo L., Verrotti A., Mastropasqua L., Chiarelli F., Della Loggia G., Morgese G., Gallenga P. E.: Colour and achromatic perimetry in diabetic children without retinopathy 247
Meynial-Denis D., Foucat L., Mignon M., Chavaroux A., Prugnand J., Bayle G., Ronou J. P., Arnal M.: Despite similar rates of alanine release, fasting and diabetes affect de novo alanine synthesis differently 606
Murakami K., Ishibashi S., Yoshida Y., Yamada N., Akanuma Y.: Lipoprotein(a) as a coronary risk factor in Japanese patients with Type II (non-insulin-dependent) diabetes mellitus. Relation with apolipoprotein(a) phenotypes 1397
Odawara M., Sasaki K., Tachi Y., Yamashita K.: Endothelial nitric oxide synthase gene polymorphism and coronary heart disease in Japanese NIDDM 365
Paquis-Flucklinger V., Vialettes B., Vague P., Canivet B., Hieronimus S., Oliver C., Pellissier J. F., Saunières A., Desnuelle C.: Importance of searching for mtDNA defects in patients with diabetes and hearing deficit 740
Rumberger J., Gottschalk W., Yarnall D., Nguyen Q., Burns D., Durcan M.: Identification of genetic markers to 20 NIDDM candidate genes by radiation hybrid analysis 366
Sakane N., Yoshida T., Yoshioka K., Nakamura Y., Umekawa T., Kogure A., Takakura Y., Kondo M.: Trp64Arg mutation of β_3-adrenoceptor gene is associated with diabetic nephropathy in Type II diabetes mellitus 1533
Sharp P. S., Al-Mrayat M., Valabhji J., Kearney T. M., Wright D.: Serum levels of vascular endothelial growth factor in diabetic subjects: the relationship with blood pressure 984
Søvik O., Njølstad P., Følling I., Sagen J., Cockburn B. N., Bell G. I.: Hyperexcitability to sulphonylurea in MODY 3 607
Sundkvist G., Tydén G., Karlsson F. A., Bolinder J.: Islet autoimmunity before and after pancreas transplantation in patients with Type I diabetes mellitus 1532
Wasada T., Katsumori K., Watanabe C., Kawahara R., Iwamoto Y.: Insulin sensitivity is inversely correlated with plasma cholesteryl ester transfer protein (CETP) 1251
Wilson C., Nelson R., Nicolson M., Pratley R.: Plasma leptin concentrations: no difference between diabetic Pima Indians with and without nephropathy 861
Winkler G., Dworak O., Salamon F., Salamon D., Speer G., Cseh K.: Increased interleukin-12 plasma concentrations in both, insulin-dependent and non-insulin-dependent diabetes mellitus 488
Winkler G., Salamon F., Salamon D., Speer G., Simon K., Cseh K.: Elevated serum tumour necrosis factor-alpha levels can contribute to the insulin resistance in Type II (non-insulin-dependent) diabetes and in obesity 860
Yamada K., Yuan X., Ishiyama S., Nonaka K.: Glucose tolerance in Japanese subjects with S20G mutation of the amylin gene 125

EASD News section

1, 17, 19, 23, 25, 27, 29, 31, 49, 57

Erratum

489, 608

Author index

1540

Subject index

1545

Contents of Volume 41

I–IX